

Fluctuation Relations for Anomalous Stochastic Dynamics

Aleksei V. Chechkin^{1,2}, Peter Dieterich³, Rainer Klages^{2,4}

1 Institute for Theoretical Physics, Kharkov, Ukraine

2 Max Planck Institute for the Physics of Complex Systems, Dresden, Germany

3 Institute for Physiology, Technical University of Dresden, Germany

4 Queen Mary University of London, School of Mathematical Sciences

Probability, Non-Local Operators and Applications
University of Sussex, 2 June 2016



Outline

- **Transient fluctuation relations (TFRs):**
motivation and warm-up
- **Correlated Gaussian dynamics:**
check TFRs for *generalized Langevin dynamics*
- **Non-Gaussian dynamics:**
check TFRs for *time-fractional Fokker-Planck equations*
- **Relations to experiments:**
glassy dynamics and biological cell migration

Motivation: Fluctuation relations

Consider a (classical) particle system evolving from some initial state into a nonequilibrium steady state.

Measure the probability distribution $\rho(\xi_t)$ of entropy production ξ_t during time t :

$$\ln \frac{\rho(\xi_t)}{\rho(-\xi_t)} = \xi_t$$

Transient Fluctuation Relation (TFR)

Evans, Cohen, Morriss (1993); Gallavotti, Cohen (1995)

why important? of *very general validity* and

- 1 generalizes the **Second Law** to small systems in noneq.
- 2 connection with **fluctuation dissipation relations**
- 3 can be checked in **experiments** (Wang et al., 2002)

Fluctuation relation for Langevin dynamics

warm-up: check TFR for the overdamped **Langevin equation**

$$\dot{x} = F + \zeta(t) \quad (\text{set all irrelevant constants to 1})$$

with **constant field** F and Gaussian white noise $\zeta(t)$.

entropy production ξ_t is equal to (mechanical) **work** $W_t = Fx(t)$
with $\rho(W_t) = F^{-1} \varrho(x, t)$; remains to solve the corresponding
Fokker-Planck equation for initial condition $x(0) = 0$:

the position pdf is Gaussian,

$$\varrho(x, t) = \frac{1}{\sqrt{2\pi\sigma_x^2}} \exp\left(-\frac{(x - \langle x \rangle)^2}{2\sigma_x^2}\right)$$

straightforward:

$$\text{(work) TFR holds if } \langle x \rangle = F\sigma_x^2/2$$

and \exists **fluctuation-dissipation relation 1 (FDR1) \Rightarrow TFR**

see, e.g., **van Zon, Cohen, PRE (2003)**

Gaussian stochastic dynamics

goal: check TFR for **Gaussian stochastic processes** defined by the (overdamped) **generalized Langevin equation**

$$\int_0^t dt' \dot{x}(t') K(t-t') = F + \zeta(t)$$

e.g., **Kubo (1965)**

with **Gaussian noise** $\zeta(t)$ and **memory kernel** $K(t)$

This dynamics can generate **anomalous diffusion**:

$$\sigma_x^2 \sim t^\alpha \text{ with } \alpha \neq 1 (t \rightarrow \infty)$$

TFR for correlated internal Gaussian noise

consider two generic cases:

1. **internal Gaussian noise** defined by the **FDR2**,

$$\langle \zeta(t)\zeta(t') \rangle \sim K(t-t'),$$

with **non-Markovian (correlated) noise**; e.g., $K(t) \sim t^{-\beta}$

solving the corresponding generalized Langevin equation in Laplace space yields

$$\text{FDR2} \Rightarrow \text{'FDR1'}$$

and since $\rho(W_t) \sim \varrho(x, t)$ is Gaussian

$$\text{'FDR1'} \Rightarrow \text{TFR}$$


for correlated internal Gaussian noise \exists TFR

Correlated external Gaussian noise


2. external Gaussian noise for which there is **no FDR2**, modeled by the (overdamped) generalized Langevin equation

$$\dot{x} = F + \zeta(t)$$

consider two types of **Gaussian noise correlated** by $g(\tau) = \langle \zeta(t)\zeta(t') \rangle_{\tau=t-t'} \sim (\Delta/\tau)^\beta$ for $\tau > \Delta$, $\beta > 0$:



persistent



anti-persistent

it is $\langle x \rangle = Ft$ and $\sigma_x^2 = 2 \int_0^t d\tau (t - \tau)g(\tau)$

Results: TFRs for correlated external Gaussian noise


σ_X^2 and the **fluctuation ratio** $R(W_t) = \ln \frac{\rho(W_t)}{\rho(-W_t)}$ for $t \gg \Delta$ and $g(\tau) = \langle \zeta(t)\zeta(t') \rangle_{\tau=t-t'} \sim (\Delta/\tau)^\beta$:

β	persistent		antipersistent *	
	σ_X^2	$R(W_t)$	σ_X^2	$R(W_t)$
$0 < \beta < 1$	$\sim t^{2-\beta}$	$\sim \frac{W_t}{t^{1-\beta}}$	regime does not exist	
$\beta = 1$	$\sim t \ln\left(\frac{t}{\Delta}\right)$	$\sim \frac{W_t}{\ln\left(\frac{t}{\Delta}\right)}$		
$1 < \beta < 2$			$\sim t^{2-\beta}$	$\sim t^{\beta-1} W_t$
$\beta = 2$	$\sim 2Dt$	$\sim \frac{W_t}{D}$	$\sim \ln(t/\Delta)$	$\sim \frac{t}{\ln\left(\frac{t}{\Delta}\right)} W_t$
$2 < \beta < \infty$			$= \text{const.}$	$\sim t W_t$

* antipersistence for $\int_0^\infty d\tau g(\tau) > 0$ yields **normal diffusion** with **generalized TFR**; above antipersistence for $\int_0^\infty d\tau g(\tau) = 0$

FDR and TFR

relation between **TFR** and **FDR I,II** for **correlated Gaussian stochastic dynamics**: ('normal FR' = conventional TFR)



in particular:

$$\boxed{\text{FDR2} \Rightarrow \text{FDR1} \Rightarrow \text{TFR}}$$

$$\boxed{\nexists \text{TFR} \Rightarrow \nexists \text{FDR2}}$$

Modeling non-Gaussian dynamics

- start again from overdamped Langevin equation $\dot{x} = F + \zeta(t)$, but here with **non-Gaussian power law correlated noise**

$$g(\tau) = \langle \zeta(t)\zeta(t') \rangle_{\tau=t-t'} \sim (K_\alpha/\tau)^{2-\alpha}, \quad 1 < \alpha < 2$$

- ‘motivates’ the **non-Markovian Fokker-Planck equation**

$$\text{type A: } \frac{\partial \varrho_A(x,t)}{\partial t} = -\frac{\partial}{\partial x} \left[F - K_\alpha D_t^{1-\alpha} \frac{\partial}{\partial x} \right] \varrho_A(x,t)$$

with **Riemann-Liouville fractional derivative** $D_t^{1-\alpha}$ (Balescu, 1997)

- two *formally similar* types derived from CTRW theory, for $0 < \alpha < 1$:

$$\text{type B: } \frac{\partial \varrho_B(x,t)}{\partial t} = -\frac{\partial}{\partial x} \left[F - K_\alpha D_t^{1-\alpha} \frac{\partial}{\partial x} \right] \varrho_B(x,t)$$

$$\text{type C: } \frac{\partial \varrho_C(x,t)}{\partial t} = -\frac{\partial}{\partial x} \left[F D_t^{1-\alpha} - K_\alpha D_t^{1-\alpha} \frac{\partial}{\partial x} \right] \varrho_C(x,t)$$

They model a *very different* class of stochastic process!

Properties of non-Gaussian dynamics

Riemann-Liouville fractional derivative defined by

$$\frac{\partial^\gamma \varrho}{\partial t^\gamma} := \begin{cases} \frac{\partial^m \varrho}{\partial t^m} & , \quad \gamma = m \\ \frac{\partial^m}{\partial t^m} \left[\frac{1}{\Gamma(m-\gamma)} \int_0^t dt' \frac{\varrho(t')}{(t-t')^{\gamma+1-m}} \right] & , \quad m-1 < \gamma < m \end{cases}$$


with $m \in \mathbb{N}$; power law inherited from correlation decay.

two important properties:


- **FDR1:** **exists** for type C but **not** for A and B
- **mean square displacement:**
 - type A: **superdiffusive**, $\sigma_x^2 \sim t^\alpha$, $1 < \alpha < 2$
 - type B: **subdiffusive**, $\sigma_x^2 \sim t^\alpha$, $0 < \alpha < 1$
 - type C: **sub-** or **superdiffusive**, $\sigma_x^2 \sim t^{2\alpha}$, $0 < \alpha < 1$
- **position pdfs:** can be calculated **approx. analytically** for A, B, only **numerically** for C

Probability distributions and fluctuation relations

• PDFs:




• TFRs:



$$R(W_t) = \log \frac{\rho(W_t)}{\rho(-W_t)} \sim \begin{cases} c_\alpha W_t, & W_t \rightarrow 0 \\ t^{(2\alpha-2)/(2-\alpha)} W_t^{\alpha/(2-\alpha)}, & W_t \rightarrow \infty \end{cases}$$

Relations to experiments: glassy dynamics

example 1: computer simulations for a **binary Lennard-Jones mixture** below the glass transition




Crisanti, Ritort, PRL (2013)

- again: $R(W_t) = \ln \frac{\rho(W_t)}{\rho(-W_t)} = \mathbf{f}_\beta(\mathbf{t}) W_t$; cp. with TFR type B
- similar results for other glassy systems (Sellitto, PRE, 2009)


Cell migration without and with chemotaxis

example 2: single MDCKF cell crawling on a substrate; trajectory recorded with a video camera



Dieterich et al., PNAS, 2008


new experiments on murine neutrophils under chemotaxis:




Dieterich et al. (2013)

Anomalous fluctuation relation for cell migration

experim. results: position pdfs $\rho(x, t)$ are **Gaussian**



fluctuation ratio $R(W_t)$ is **time dependent**



$\langle x(t) \rangle \sim t$ and $\sigma_x^2 \sim t^{2-\beta}$ with $0 < \beta < 1$: \nexists FDR1 and

$$R(W_t) = \ln \frac{\rho(W_t)}{\rho(-W_t)} = \frac{W_t}{t^{1-\beta}}$$

data matches to analytical results for persistent correlations

Summary

- TFR tested for two generic cases of **correlated Gaussian stochastic dynamics**:
 - 1 **internal noise**:
FDR2 implies the validity of the 'normal' work TFR
 - 2 **external noise**:
FDR2 is broken; sub-classes of **persistent** and **anti-persistent noise** yield both **anomalous TFRs**
- TFR tested for three cases of **non-Gaussian dynamics**:
breaking FDR1 implies again **anomalous TFRs**
- anomalous TFRs appear to be important for **glassy aging dynamics**: cf. computer simulations on various glassy models and experiments on ('gelly') cell migration

References

- P.Dieterich, RK, A.V. Chechkin, NJP **17**, 075004 (2015)
- A.V. Chechkin, F.Lenz, RK, J. Stat. Mech. L11001 (2012)
- A.V. Chechkin, RK, J. Stat. Mech. L03002 (2009)

